

Health Care for Children, Pregnant & Breastfeeding women, and Geriatric patients

Prepared by

Dr. Ayesha Fatima

Assistant Professor

Dept of Pharmacy Practice

The Pediatric Health Ecosystem: First-line connective tissue for child development.



Pharmacists are not just dispensers; they are the proactive educators, safety monitors, and primary clinical touchpoints ensuring these foundational pillars remain intact from birth through adolescence.

Health Promotion in Children (Birth to Adolescence)

1. From birth to adolescence, children require continuous support from parents, caregivers, and healthcare professionals to achieve optimal physical, mental, emotional, and social development.
2. The Pediatric age group generally includes children from 0–16 years.
3. During different stages of growth, children experience various health problems and are at risk of infections, nutritional deficiencies, developmental disorders, and accidents.
4. Based on age, children are classified into neonates, infants, children, and adolescents.
5. Health promotion during childhood aims to prevent disease, promote healthy growth, and establish lifelong healthy behaviours.

The major components of health promotion in children include:

0 Years

14 Years

Continuous clinical support across early development prevents disease and establishes lifelong behaviors.



Immunization

Strengthening the immune system against infectious diseases.



Nutrition

Fueling physical and brain development with balanced energy.



Oral Hygiene

Preventing childhood dental caries through active daily care.



Healthy Habits

Promoting physical and mental well-being via sleep, diet, and activity.

1. Immunization


- ❖ Immunization is the process by which an individual's immune system is strengthened against infectious diseases by exposing it to an immunogen, usually through vaccination.
- ❖ Vaccines stimulate the body's immune system to produce protective antibodies, thereby preventing infections or reducing disease severity.
- ❖ Immunization is one of the most effective methods of preventing childhood illnesses and reducing child mortality.
- ❖ Children should receive vaccines according to the national immunization schedule from birth through adolescence.

Vaccines introduce controlled immunogens to stimulate long-term protection.



Step 1: Introduction

Introduction of
Immunogen.



Step 2: Antibody Production


Immune system
produces protective
antibodies.



Step 3: Long-term Shield

Prevention of infection
and reduction of
disease severity.

The 5-Year Childhood Immunization Schedule



General Childhood Immunization Schedule

1. **At birth:** First dose of Hepatitis B vaccine.
2. **At 1 month:** Second dose of Hepatitis B vaccine.
3. **At 6–8 weeks:** First dose of DPT (Diphtheria, Pertussis, Tetanus) and Polio vaccines.
4. **At 3 months:** Second dose of DPT and Polio vaccines.
5. **At 4 months:** Third dose of DPT and Polio vaccines.
6. **At 12–15 months (around 13 months):** MMR (Measles, Mumps, and Rubella) vaccine along with Hepatitis B booster if indicated.
7. **At 4–5 years:** Booster doses of DPT and MMR vaccines.

Timely vaccination provides long-term protection against many serious infectious diseases.

Clinical Playbook: The Pharmacist's Role in Immunization



- ✔ **Educate & Advise:** Counsel parents on vaccine schedules, safety, and managing possible adverse effects.
- ✔ **Maintain Access:** Ensure adequate stock of recommended vaccines.
- ✔ **Public Health Activation:** Support vaccination drives during outbreaks (e.g., influenza, H1N1) and participate in school health programs.
- ✔ **Administer & Refer:** Act as immunization facilitators (administering where permitted by law) and refer to physicians when required.

Role of Community Pharmacists in Immunization

Community pharmacists play an important role in promoting childhood immunization by:


1. Serving as the first point of healthcare contact for many families.
2. Maintaining adequate stocks of recommended vaccines.
3. Educating parents about vaccine schedules and their importance.
4. Advising parents regarding vaccine safety and possible adverse effects.
5. Referring children to physicians or immunization centers when required.
6. Participating in public awareness campaigns and school health education programs.
7. Supporting vaccination drives during disease outbreaks such as influenza and H1N1.
8. Acting as immunization facilitators and, where permitted by law, administering vaccines.

2. Nutrition

- Proper nutrition is essential for normal physical growth, brain development, immune function, and overall health throughout childhood. Establishing healthy eating habits early in life provides lifelong health benefits and reduces the risk of chronic diseases.
- A balanced diet should supply adequate amounts of:
 1. Carbohydrates
 2. Proteins
 3. Healthy fats
 4. Vitamins
 5. Minerals
 6. Dietary fiber
 7. Water
- Children should be encouraged to consume fruits, vegetables, whole grains, milk, dairy products, eggs, fish, lean meat, and nuts while limiting foods high in fat, sugar, and salt.

Caloric requirements scale linearly to fuel rapid pediatric growth

$$\text{Daily calories} = 1000 + (100 \times \text{Age in years})$$



Note: Exact requirements vary depending on body size, growth rate, and daily physical activity level. Diet must balance carbs, proteins, fats, vitamins, minerals, fiber, and water.

Daily Calorie Requirement

- Children require adequate calories to support growth, development, and daily physical activities.
- A simple method for estimating calorie requirements for children up to five years of age is:

$$\text{Daily calories} = 1000 + (100 \times \text{Age in years})$$

- For example:
 1. 1-year-old child: approximately **1100 kcal/day**
 2. 2-year-old child: approximately **1200 kcal/day**
 3. 3-year-old child: approximately **1300 kcal/day**
- Calorie requirements vary depending on body size, growth rate, and activity level.

The Weaning Window

At six months, expanding nutritional needs require the gradual introduction of solid foods.



Guidance:

- **Start Small:** Begin with 1 teaspoon, 1-2 times daily.
- **Pacing:** Introduce one new food at a time to monitor for allergic reactions.
- **Texture:** Ensure foods are soft, mashed, or pureed until chewing develops.

Navigating the three primary modalities of infant nutrition.



Breast Milk

The Preferred Method

- **Timeline:** 0-6 months (exclusive).
- **Benefit:** Complete nutrition & infection protection.



Infant Formula

The Alternative

- **Timeline:** When breastfeeding is insufficient.
- **Benefit:** Provides essential nutrients & prevents anemia.
- **Requirement:** Must be iron-fortified; no extra vitamins needed.
- **Warning:** Never use homemade substitutes.




Solid Foods

The Complement

- **Timeline:** 6+ months.
- **Benefit:** Meets expanding nutritional requirements.
- **Foods:** Iron-fortified cereals, pureed veg, fruit, and meat.

The Caries Risk Scale

Dental caries are highly prevalent but entirely preventable through balanced oral hygiene.



Infant Feeding

1. Breastfeeding

Breastfeeding is the preferred method of infant feeding. Exclusive breastfeeding is recommended during the first **six months** of life because breast milk provides complete nutrition and protects against infections.


2. Infant Formula

When breastfeeding is not possible or is insufficient, commercially prepared **iron-fortified infant formula** is recommended.

Infant formulas:

1. Provide essential nutrients.
2. Help prevent iron-deficiency anemia.
3. Should be used under medical guidance.
4. Should not be replaced with homemade formulas because they may be nutritionally inadequate or unsafe.
5. Formula-fed infants generally do not require additional vitamin supplementation.

The Formula Preparation Safety Lifecycle



Preparation of Infant Formula

1. Proper preparation is essential to ensure safety.
2. Follow the manufacturer's mixing instructions exactly.
3. Use boiled and cooled water, especially during the first three months.
4. Sterilize feeding bottles and utensils before use.
5. Check the expiry date before preparing formula.
6. Store formula according to the manufacturer's instructions.
7. Discard leftover formula after feeding.
8. Do not heat formula directly on the stove or in a microwave.
9. Warm bottles by placing them in lukewarm water.
10. Never leave a baby unattended with a propped bottle because of the risk of choking.

Introduction of Solid Foods

Breast milk or formula alone becomes insufficient to meet all nutritional requirements after six months of age.

- Current recommendations include:
 1. Exclusive breastfeeding for the first **6 months**.
 2. Introduce complementary foods at **6 months** while continuing breastfeeding.
 3. Suitable first foods include:
 1. Iron-fortified cereals
 2. Pureed vegetables
 3. Pureed fruits
 4. Mashed pulses
 5. Pureed chicken or meat
 6. Cottage cheese
 4. Initially: Begin with one teaspoon once or twice daily.
 5. Gradually increase both quantity and variety.
 6. Introduce one new food at a time.
 7. Observe for allergic reactions.
 8. Ensure homemade foods are soft, mashed, or pureed until chewing develops.
 9. Parents should consult healthcare professionals before introducing complementary foods.

Clinical Playbook: The Pharmacist's Role in Nutrition



Inventory Provision:

Stock appropriate infant formulas and baby foods.

Feeding Counsel:

Advise parents on breastfeeding techniques, formula feeding, and the introduction of age-appropriate complementary foods.

Safety Demonstration:

Physically demonstrate correct formula preparation and bottle sterilization techniques to caregivers.

Nutritional Education:

Educate families on balanced diets and the prevention of pediatric nutritional deficiencies.

Role of Pharmacists in Child Nutrition

- Pharmacists contribute to child nutrition by:
 1. Stocking appropriate infant formulas and baby foods.
 2. Counseling parents on breastfeeding and formula feeding.
 3. Demonstrating correct formula preparation.
 4. Explaining bottle sterilization techniques.
 5. Advising on age-appropriate complementary feeding.
 6. Educating parents about balanced nutrition and prevention of nutritional deficiencies.

3. Oral Hygiene

Good oral health is essential for overall health and well-being. Dental caries (tooth decay) is one of the most common childhood diseases but is largely preventable through proper oral hygiene and healthy dietary habits.

Risk factors include:

1. Frequent consumption of sugary foods and drinks.
2. Poor brushing habits.
3. Inadequate fluoride exposure.
4. Poor dental care.

Children should:

1. Brush teeth twice daily with fluoride toothpaste.
2. Reduce sugary snacks and beverages.
3. Visit a dentist regularly.
4. Maintain proper oral hygiene from the eruption of the first tooth.

Clinical Playbook: The Pharmacist's Role in Oral Health



Product Recommendation: Recommend fluoride toothpaste, fluoride supplements, and sugar-free medicines/lozenges.

Habit Modification: Encourage transitioning to drinking cups over prolonged bottle feeding. Guidance on oral appliance hygiene if necessary.

Triage & Relief: Provide analgesics for temporary dental pain relief.

Clinical Referral: Refer patients to dentists for definitive treatment, persistent mouth ulcers, or oral lesions. Participate in dental awareness camps.

Role of Community Pharmacists in Oral Health

Community pharmacists promote oral health by:

1. Educating parents about proper oral hygiene.
2. Recommending fluoride toothpaste and fluoride supplements when appropriate.
3. Advising on sugar-free medicines and lozenges.
4. Encouraging the use of drinking cups instead of prolonged bottle feeding.
5. Providing guidance on denture hygiene when necessary.
6. Referring patients with persistent mouth ulcers or oral lesions to dentists.
7. Participating in oral health awareness programs and dental camps.
8. Providing analgesics for temporary relief of dental pain and referring patients for definitive treatment

Establishing physical and mental well-being through daily habit tracking.



Handwashing

6 crucial touchpoints: Before eating, after toilet, nappies, coughing/blowing nose, outdoors, handling toys/animals.



Physical Activity

Moderate-intensity daily activity (ages 2+). Limit sedentary screens. Ensure hydration.



Balanced Diet

Regular meals balancing fruits, veg, proteins, grains. Multivitamins only under medical advice for poor intake.



Adequate Sleep



Regular sleep schedules are mandatory for physical growth, learning, memory, and emotional behavior.

4. Healthy Habits in Children

- Healthy habits established during early childhood often continue throughout life.
- Parents and caregivers should consistently encourage behaviors that promote physical and mental well-being.

a. Good Hand Washing

Proper hand washing is one of the simplest and most effective methods of preventing infectious diseases.

Children should wash their hands:

1. Before eating.
2. After using the toilet.
3. After changing nappies.
4. After coughing or blowing the nose.
5. After playing outdoors.
6. After handling toys or animals.
7. Hand washing with soap and clean water removes dirt and disease-causing microorganisms.

b. Adequate Sleep

Adequate sleep is essential for:

1. Physical growth.
2. Brain development.
3. Learning and memory.
4. Emotional well-being.
5. Good concentration and behavior.

Sleep requirements vary according to age, but children should maintain regular sleep schedules.

c. Healthy Eating

Children should consume a balanced diet containing:

1. Fruits and Vegetables
2. Milk and dairy products
3. Eggs
4. Fish
5. Chicken
6. Meat
7. Whole grains
8. Nuts

Meals should be regular, nutritious, and balanced. Multivitamin supplementation may be considered for children with poor dietary intake under medical advice.

d. Regular Physical Activity

Regular exercise promotes:

1. Healthy growth.
2. Strong bones and muscles.
3. Cardiovascular fitness.
4. Healthy body weight.
5. Better mental health.

Children aged 2 years and older should participate in at least 30 minutes of enjoyable moderate-intensity physical activity every day.

Sedentary activities such as excessive television viewing, mobile phone use, and video games should be limited.

Adequate water intake should also be encouraged during physical activity.

Health Promotion and Care for Pregnant Women and Breastfeeding Mothers

1. Pregnancy is a critical period that requires proper medical care to ensure the health of both the mother and the developing fetus.
2. Good prenatal (antenatal) care reduces maternal complications and fetal mortality.
3. The main goal of prenatal care is to ensure the safe delivery of a healthy baby.

Antenatal (Prenatal) Care

1. Early Confirmation of Pregnancy

1. Pregnancy can be confirmed using home pregnancy test kits.
2. A gynecologist can detect pregnancy as early as 8–9 days after fertilization.
3. Early diagnosis allows timely initiation of antenatal care.

2. First Antenatal Visit

During the first visit, the healthcare provider performs a complete assessment.

• Maternal Assessment

1. Measurement of: Blood pressure , Height, Weight
2. Assessment of overall health status.
3. Review of medical and obstetric history.
4. Laboratory Investigations

1. Blood tests

1. Blood group and Rh typing
2. Hemoglobin estimation
3. Screening for infections
4. Screening for diseases such as diabetes and hepatitis (as indicated)

2. Urine tests

1. Sugar
2. Protein
3. Urinary tract infection

Pelvic Examination

1. Evaluation of the uterus (womb) size.
2. Assessment of reproductive organs.
3. Confirmation of pregnancy progression.

Estimation of Gestational Age

1. Calculated using the Last Menstrual Period (LMP).
2. Expressed in completed weeks of gestation.
3. Helps determine:
 1. Expected Date of Delivery (EDD)
 2. Growth of the fetus
 3. Schedule of antenatal visits.

Health Promotion During Pregnancy

1. Balanced Nutrition

1. Consume a balanced diet rich in:
 1. Proteins
 2. Iron
 3. Calcium
 4. Folic acid
 5. Vitamins
2. Drink adequate fluids.
3. Maintain appropriate weight gain.
4. Good nutrition promotes healthy fetal growth and maternal well-being.

2. Adequate Rest and Sleep

1. Ensure sufficient sleep every day.
2. Avoid excessive physical and mental stress.
3. Proper rest improves maternal health and fetal development.

3. Regular Physical Activity

1. Perform moderate exercise as advised by the doctor.
2. Benefits include:
 1. Better circulation
 2. Reduced back pain
 3. Easier labor
 4. Improved physical fitness

5. Regular Antenatal Check-ups

Regular visits help monitor:

1. Maternal blood pressure
2. Weight gain
3. Fetal growth
4. Fetal heartbeat
5. Early detection of pregnancy complications

4. Avoid Harmful Habits

Pregnant women should avoid:

1. Smoking
2. Alcohol consumption
3. Tobacco products
4. Recreational drugs
5. Self-medication
6. Unnecessary Over-the-Counter

These substances may cause:

1. Birth defects
2. Poor fetal growth
3. Premature birth
4. Miscarriage

Importance of Prenatal Care

1. Detects pregnancy complications early.
2. Reduces maternal mortality.
3. Decreases fetal mortality.
4. Promotes healthy fetal development.
5. Improves maternal health.
6. Ensures safe pregnancy and delivery.
7. Provides individualized care according to maternal needs.

Breastfeeding After Delivery

Importance of Breastfeeding

1. Breastfeeding benefits both the mother and the baby.
2. Breast milk is the ideal nutrition for infants.

Colostrum

1. Colostrum is the first milk produced after childbirth.
2. Thick, yellowish, and highly nutritious.
3. Rich in:
 1. Antibodies (especially IgA)
 2. Proteins
 3. Immune cells
4. Provides passive immunity.
5. Protects the newborn from infections.

Benefits of Breastfeeding for the Baby

1. Provides complete nutrition.
2. Easily digested.
3. Strengthens the immune system.
4. Reduces infections (diarrhea, respiratory infections).
5. Promotes healthy growth and development.
6. Enhances emotional bonding with the mother.

Benefits of Breastfeeding for the Mother

1. Promotes mother–baby bonding.
2. Increases maternal confidence and self-esteem.
3. Helps uterine contraction after delivery.
4. Reduces postpartum bleeding.
5. Lowers the risk of breast cancer.
6. May reduce the risk of ovarian cancer.
7. Helps in weight reduction after pregnancy.

Role of the Pharmacist in Counseling Pregnant Women

The pharmacist plays an important role in promoting maternal and fetal health.

During Pregnancy

1. Dispense safe and quality medicines.
2. Educate women about the safe use of medications during pregnancy.
3. Counsel against self-medication.
4. Advise against unnecessary OTC drugs.
5. Explain the importance of:
 1. Balanced diet
 2. Adequate rest
 3. Proper sleep
 4. Regular exercise
6. Educate about avoiding:
 1. Smoking
 2. Alcohol
 3. Tobacco
 4. Harmful substances
7. Reinforce adherence to prescribed medications (e.g., iron, folic acid, calcium).
8. Encourage regular antenatal check-ups.

After Delivery

1. Educate mothers about the importance of exclusive breastfeeding for the first 6 months.
2. Explain the benefits of colostrum.
3. Counsel on correct breastfeeding techniques.
4. Provide advice regarding infant formula only when breastfeeding is not possible.
5. Inform mothers that some medicines are excreted into breast milk.
6. Advise consultation with healthcare professionals before taking any medicine while breastfeeding.
7. Monitor for medicines that may affect the infant through breast milk.

Health Promotion and Health Care of Geriatric Patients

- Geriatrics is the branch of medicine concerned with the clinical, preventive, rehabilitative, and social aspects of health and illness in older adults.
- Elderly generally refers to individuals 60 years of age and above.
- India is undergoing a demographic transition, resulting in a rapid increase in the elderly population.
- Average life expectancy has increased significantly and is expected to improve further.
- India has one of the largest elderly populations in the world, and by 2050, elderly individuals are expected to constitute a major proportion of the global aging population.
- Increased longevity has created new challenges in maintaining health, independence, and quality of life among older adults.
- Health promotion in geriatrics refers to helping older adults maintain good physical, mental, and social health, prevent diseases, delay disability, and improve their quality of life.

The Demographic Transition and the Ageing Reality



2050 Projection: India is expected to harbor one of the largest elderly (60+ years) populations in the world, constituting a major proportion of the global aging demographic.

The Core Challenge

Average life expectancy has increased significantly. However, this increased longevity creates immediate, critical challenges in maintaining health, independence, and quality of life.


The Required Pivot

Health promotion must shift from merely increasing lifespan to actively ensuring healthy ageing.

Need for Health Promotion for Geriatrics

- Increasing life expectancy has resulted in:
 1. Increased prevalence of chronic diseases
 2. Functional disability
 3. Polypharmacy
 4. Greater healthcare utilization
 5. Reduced quality of life
 6. Increased healthcare costs
- Therefore, health promotion focuses on healthy ageing rather than merely increasing lifespan.

The Reality of Geriatric Healthcare



The Treatment Burden

These coexisting illnesses necessitate **long-term therapy**.

Patients must navigate multiple physicians and frequent hospital visits.

Result: Older adults are the largest consumers of medicines, frequently taking 5 or more prescription drugs daily alongside OTCs and supplements.

Burden of Non-Communicable Diseases (NCDs)


- One of the major determinants of health in old age is the presence of chronic non-communicable diseases
- Common NCDs include:
 1. Hypertension
 2. Diabetes mellitus
 3. Coronary heart disease
 4. Stroke
 5. Chronic obstructive pulmonary disease (COPD)
 6. Cancer
 7. Arthritis
 8. Osteoporosis
- These diseases are responsible for:
 1. Disability
 2. Functional dependence
 3. Reduced quality of life
 4. Increased mortality

Life-Course Approach

- WHO recommends a life-course approach for preventing NCDs.
- Risk factors for chronic diseases begin much earlier than old age.
- They may originate during:
 1. Pregnancy (maternal health)
 2. Childhood
 3. Adolescence
 4. Adult life

Therefore, healthy lifestyle practices throughout life help delay or prevent diseases in old age.

The Life-Course Trajectory of Chronic Disease



The WHO Life-Course Approach

Geriatric health is a compounding process. Healthy lifestyle practices must be sustained throughout life to delay or prevent disease in old age.

Major Modifiable Risk Factors

- The major lifestyle factors responsible for NCDs include:
 1. Physical inactivity
 2. Unhealthy diet
 1. High calories
 2. High sugar
 3. High saturated fat
 4. Excess salt
 5. Low intake of fruits and vegetables
 3. Tobacco use
 4. Alcohol consumption
 5. Obesity
 6. Sedentary lifestyle

Relationship Between Risk Factors and Diseases

Risk Factor	Diseases
Tobacco use	Cancer, COPD, Heart disease, Stroke
Physical inactivity	Diabetes, Coronary heart disease, Stroke
Unhealthy diet	Obesity, Diabetes, Hypertension, Heart disease
Alcohol	Liver disease, Hypertension, Stroke, Certain cancers

Mapping Modifiable Risks to NCD Outcomes

	Cancer	COPD	Heart Disease	Stroke	Diabetes	Hypertension
Tobacco Use	✓	✓	✓	✓		
Physical Inactivity			✓	✓	✓	
Unhealthy Diet (High Sugar/Fat/Salt)			✓		✓	✓
Alcohol Consumption	✓			✓		✓

These modifiable NCDs are the primary drivers of functional dependence, reduced quality of life, and escalating healthcare expenditure in geriatric patients.

Health Transition in India

- India is currently experiencing:
 1. Rapid urbanization
 2. Industrialization
 3. Lifestyle changes
 4. Population ageing
- As a result:
 - Non-communicable diseases have become the leading cause of illness.
 - India is known as the "**Diabetes Capital of the World.**"
 - India carries approximately one-fourth of the world's burden of coronary heart disease.
 - Indians develop diabetes and coronary artery disease at a younger age than many developed countries.
 - Survivors therefore live with disability for a longer period.

Expansion of Morbidity

Although life expectancy is increasing, many elderly spend more years living with chronic illness.

This phenomenon is known as:

- **Expansion of Morbidity**

1. Instead of compressing illness into the last few years of life, chronic diseases now extend over many years.
2. Therefore, emphasis should be on:
3. Prevention
4. Early diagnosis
5. Lifestyle modification
6. Regular screening

The Expansion of Morbidity

Historical Trajectory: Shorter Lifespan



Current Reality: Extended Lifespan



Context: India's Health Transition

- ✓ Known as the Diabetes Capital of the World, carrying ~25% of the global coronary heart disease burden.
- ✓ Populations are developing diabetes and coronary artery disease at significantly younger ages.
- ✓ Result: Survivors live with functional dependence and disability for far longer periods.

Goals of Health Promotion in Geriatrics

1. Promote healthy ageing
2. Prevent chronic diseases
3. Delay disability
4. Maintain independence
5. Improve quality of life
6. Reduce hospitalization
7. Reduce healthcare expenditure
8. Promote active ageing

Medication Use in Older Adults

- Older adults are the **largest consumers of medicines**, including both prescription and over-the-counter (OTC) drugs.
- Reasons include:
 1. Multiple chronic diseases
 2. Long-term therapy
 3. Multiple physicians
 4. Frequent hospital visits
- Older adults often take **five or more prescription medicines daily**.
- Many also consume OTC medications, vitamins, herbal medicines, and nutritional supplements.

This increases the risk of:

1. Polypharmacy
2. Drug interactions
3. Adverse drug reactions
4. Medication errors
5. Poor adherence

The Medication Perception Gap: OTC and Herbal Risks

Elderly Patient Belief

“Over-the-counter medicines, vitamins, dietary supplements, and herbal remedies are completely safe and mild.”


Clinical Reality

- ⚠ Many potent prescription medicines have transitioned to OTC.
- ⚠ They produce significant physiological effects and can cause severe interactions with existing prescriptions.
- ⚠ Herbal medicines and dietary supplements are not exempt from serious, life-threatening interactions.

Action: OTC and herbal medicines must be treated with the exact same caution as prescription therapies.

The Polypharmacy Cascade

Definition: The simultaneous use of five or more medications.



Polypharmacy

- Polypharmacy is the use of five or more medications simultaneously.
- **Consequences**
 1. Drug-drug interactions
 2. Drug-food interactions
 3. Drug-disease interactions
 4. Falls
 5. Confusion
 6. Hospitalization
 7. Increased mortality
 8. Poor medication adherence

Use of OTC Medicines

Many elderly believe OTC medicines are completely safe.

However,

1. Many prescription medicines have now become OTC medicines.
2. OTC medicines can produce significant physiological effects.
3. They may interact with prescription drugs.
4. Herbal medicines and dietary supplements may also cause serious interactions.

Therefore, OTC medicines should be used cautiously.

Barrier-Solution Matching for Medication Adherence

Physical & Cognitive Barriers

Arthritis & Limited Dexterity
Difficulty opening standard containers.

Poor Eyesight
Struggles reading small print labels.

Memory Problems
Forgetting doses or taking them incorrectly.

Complex Regimens
Multiple medications with varying schedules.

Severe Cognitive Impairment
Significant decline in cognitive function.

Practical Solutions

Easy-open medicine containers.
Utilize non-child-resistant caps.

Large-print labels, magnifying glasses, and adequate lighting.
Enhance readability.

Routine anchoring (taking meds with meals), or physically turning bottles upside down after dosing.
Create behavioral cues.

Requesting once-daily or twice-daily simplified dosing schedules.
Minimize regimen complexity.

Engaging caregivers, home health nurses, or adult day-care services.
Provide necessary support and supervision.

Common Drug-Related Problems in Elderly

- Older adults should immediately consult a healthcare professional if they experience:
 1. Dizziness
 2. Constipation
 3. Diarrhea
 4. Stomach upset
 5. Blurred vision
 6. Sleep disturbances
 7. Mood changes
 8. Skin rashes
 9. Unusual weakness
 10. Confusion
- These symptoms may indicate adverse drug reactions rather than ageing itself.

Diagnostic Trap: Ageing vs. Adverse Reactions



Symptoms Often Dismissed as Natural Ageing

- ✓ Dizziness & Unusual weakness
- ✓ Confusion & Mood changes
- ✓ Sleep disturbances
- ✓ Blurred vision
- ✓ Constipation, Diarrhea, Stomach upset
- ✓ Skin rashes

Indicators of Adverse Drug Reactions

These are not inevitable signs of getting older. These are highly indicative of dangerous, medication-related problems requiring immediate intervention by a healthcare professional.

Safe Medication Practices for Older Adults

Practice	Recommendation
Inform Your Doctor/Pharmacist	Tell them about all prescription medicines, OTC drugs, vitamins, herbal products, and allergies.
Use One Pharmacy	Get all medicines from the same pharmacy to avoid drug interactions.
Watch for Side Effects	Report dizziness, rashes, stomach upset, blurred vision, or other unusual symptoms.
Know Your Medicines	Learn the name, dose, timing, purpose, and possible side effects of each medicine.
Review Medicines Regularly	Bring all your medicines during doctor visits for review.
Follow Instructions	Take medicines exactly as prescribed and read the label carefully.
Use Reminder Aids	Use a pill box, calendar, or alarm to remember your medicines.
Avoid Self-Medication	Do not start or stop medicines without consulting your doctor or pharmacist.

The Pharmacist as the Care Anchor

Direct Clinical Interventions

- **Patient Counseling**
(dosage, adverse effects, adherence)
- **Medication Review**
(identifying polypharmacy, duplicate therapy)
- **Health Screening**
(monitoring BP, glucose, BMI)

Holistic & Community

- **Lifestyle Counseling**
(diet, sleep hygiene, stress)
- **Promoting Healthy Behaviour**
(vaccinations, exercise)
- **Community Education**
(senior citizen clubs, peer-support)

The Patient

Multidisciplinary & Extension

- **Physician Collaboration**
(discussing dose adjustments, deprescribing)
- **Home Medication Reviews**
(assessing storage, caregiver education)
- **Improving Adherence**
(managing schedules, preventing missed doses)

Role of Pharmacist in Geriatric Care

Role	Responsibilities
Patient Counseling	Educate elderly patients about their diseases, medicines, dosage, side effects, and the importance of medication adherence.
Medication Review	Review all medicines regularly to identify polypharmacy, drug interactions, duplicate therapy, and adverse drug reactions.
Health Screening	Monitor blood pressure, blood glucose, body weight, and other basic health parameters.
Medication Adherence	Encourage patients to take medicines correctly and use pill boxes, calendars, or alarms as reminders.
Lifestyle Counseling	Advise on a healthy diet, regular exercise, smoking cessation, alcohol reduction, stress management, and personal hygiene.

OTC and Herbal Medicine Guidance	Counsel patients on the safe use of over-the-counter medicines and herbal supplements to avoid harmful interactions.
Home Medication Review	Assess medicines at home for proper storage, adherence, expired drugs, and medication-related problems.
Collaboration with Physicians	Communicate with doctors regarding drug-related problems, dose adjustments, adverse effects, and therapeutic alternatives.
Health Education	Conduct health awareness programs, group counseling, and senior citizen education sessions to promote healthy ageing.
Promote Healthy Ageing	Encourage regular health check-ups, vaccination, disease prevention, and healthy lifestyle practices to improve quality of life.

The 7-Pillar Safe Medication Dashboard



1. One Pharmacy

Fill all prescriptions at a single pharmacy to detect interactions and maintain a master record.



2. Full Disclosure

Inform doctors of all OTC, herbal, and vitamin usage, plus allergies.



3. Monitor Effects

Never assume new symptoms are just old age.



4. Learn the Regimen

Know drug name, dose, frequency, duration, side effects, and food interactions.



5. Medication Review

Bring all medicines to every doctor visit to spot duplicates.



6. Exact Adherence


Read labels carefully for exact timing, dose, and food instructions.



7. Reminder Systems

Utilize pill organizers, calendars, and mobile alarms.

Synthesizing the Blueprint for Active Ageing



By addressing the life-course trajectory, dismantling the polypharmacy cascade, and elevating the multidisciplinary team, we can transform the expansion of morbidity into true, active ageing.